


 Search

- Home
 - Genomics
 - News & Information
 - Research
 - Cores
 - Resources
 - People
 - Workspaces
 - My Account
 - About Us
-
- All
 - Adenoviruses
 - Antibodies
 - Bioimages
 - mESC Lines
 - Mouse Strains
 - Genomics Studies
 - Protocols
 - Miscellaneous
 - Research Data
 - Visualization

- My Account**
 - Login
 - Create Account
-
- Resources**
 - View All (813)
 - Adenoviruses (137)
 - Antibodies (175)
 - Bioimages (67)
 - Genomics Studies (145)
 - mESC Lines (68)
 - Mouse Strains (120)
 - Miscellaneous (46)
 - Protocols (55)
 - Research Data (4)
 - Resource Tags (389)
 - Visualization (9)
-
- Research & Cores**
 - Core Facilities (5)
 - Research Highlights (5)
 - Research Networks
 - Research Objectives
-
- Information**
 - About the BCBC
 - BCBC Events
 - Branding & Logos
 - Career Opportunities
 - Health
 - NIH hESC Registry
 - Policies & Guidelines
 - Member Publications
 - Research Programs
 - Research Investigators
 - Member Directory
 - Tutorials

Comparison of endocrine enriched genes in islet beta cells versus induced beta cells - Study GBCO3549

Genomics Study Specifications

Study Name	Comparison of endocrine enriched genes in islet beta cells versus induced beta cells
Contact Name	Douglas Melton (Harvard University)
Publication	http://www.ncbi.nlm.nih.gov/pubmed/18754011
My Strategies	Return to My Strategies page
Classification	Targets and roles of transcriptional regulators; Pancreas development and growth
Links	Biomaterials Graph GEO
BCBC Release Date	February 09, 2009
Public Release Date	February 09, 2009
Citation	Zhou Q, Brown J, Kanarek A, Rajagopal J, Melton DA. In vivo reprogramming of adult pancreatic exocrine cells to beta-cells . Nature. 2008. 455:627-32
Synopsis	<div style="border: 1px solid #ccc; padding: 5px;"> <div style="display: flex; justify-content: space-between; border-bottom: 1px solid #ccc;"> Study Description Goals </div> <div style="display: flex; justify-content: space-between; border-bottom: 1px solid #ccc;"> Approaches Results Conclusions </div> <div style="display: flex; justify-content: space-between;"> Related Studies </div> <p>Endocrine enriched genes in adult islet beta cells were identified and compared with that of induced beta cells (with M3 transcription factors) in adult. The control sample is non-beta pancreatic cells. Gene expression profile comparison of 3 samples, 3 independent repeats for each sample indicate a high degree of similarity between endogenous and induced beta cells in adult mouse.</p> </div>
Platform types	Expression microarray, Expression
Platforms	Show platform Illumina mouseRef-8 v1.1 expression beadchip
Study Design Type	<ul style="list-style-type: none"> • cell_type_comparison_design • genetic_modification_design
Study Factors	Show study factors
Study Assays	Show study assays

Access to Study Data

This Study Data is publicly available to all users.

Gene List(s)

Use the following form(s) to refine the parameters and add the gene list to a strategy:

▼
Mouse induced beta cells versus non-islet pancreatic cells

Access Status

This resource is publicly viewable.

Request this Resource

Request from a repository

Primary contributor: [Melton Lab](#)

- Co-contributed by:
- [Stoeckert Lab](#)

Resource Tags

1a, Amy2, amylase, amylase 2, beta 3, Gcg, Gck, glucagon, glucokinase, Glut2, Illumina mouseRef-8 v1.1 expression beadchip, Ins2, insulin, insulin II, Isl1, ISL1 transcription factor, keratin 19, Krt19, LIM/homeodomain, locus 2 (Drosophila), Mafa, member 2, Nes, nestin, Neurod, Neurod1, Neurog3, neurogenic differentiation 1, neurogenin 3, ngn3, NK2 transcription factor related, NK6 homeobox 1, Nkx2-2, Nkx2.2, Nkx6-1, Nkx6.1, paired box gene 4, paired box gene 6, pancreas specific transcription factor, pancreatic, pancreatic and duodenal homeobox 1, pancreatic polypeptide, Pax4, Pax6, Pcsk1, Pdx1, Ppy, proprotein convertase subtilisin/kexin type 1, protein A (avian), Ptf1a, Slc2a2, solute carrier family 2 (facilitated glucose transporter), somatostatin, Sst, test, Tubb3, tubulin, Vim, vimentin, v-maf musculoaponeurotic fibrosarcoma oncogene family

Login to edit tags

[Read more about tags](#)

Resource History & Actions

Approved on Feb 09, 2009
Last modified on Aug 02, 2011

Login to edit or request an edit

Related resources

BCBC

No matching resources

Other Consortia

No matching resources

Data courtesy of [dkCOIN](#). Only public resources are displayed.

|Fold Change| Greater Than:

Confidence Level:

High Confidence All Results

For a microarray experiment a result with high confidence has a confidence level of at least 80%.

For a ChIP-chip experiment a result with high confidence has a confidence level of at least 90% and all fold changes are positive.

Reference (Denominator):

Non-islet pancreatic cells

[Find Genes](#)

▶ **Mouse whole islet cells versus non-islet pancreatic cells**

Genome Browser


There are no genome browser tracks currently available for this study.

Lists of Locations

There are no genomic location datasets currently available for this study.

Repositories

Melton Lab


 Request this resource

Stock #: *Not provided*

Availability Notes: *Not provided*

Comments

There are no comments for this entry.

 Login to add comments

